Precision Theoretical Analysis of Neutron Radiative Beta Decay

A. N. Ivanov,[1,*] R. Höllwieser,[1,2,†] N. I. Troitskaya,[1,‡] M. Wellenzohn,[1,3,§] and Ya. A. Berdnikov[4,¶]

[1] *Atominstitut, Technische Universität Wien, Stadionallee 2, A-1020 Wien, Austria*
[2] *Department of Physics, New Mexico State University, Las Cruces, New Mexico 88003, USA*
[3] *FH Campus Wien, University of Applied Sciences, Favoritenstraße 226, 1100 Wien, Austria*
[4] *Peter the Great St. Petersburg Polytechnic University, Polytechnicheskaya 29, 195251, Russian Federation*

(Dated: January 18, 2017)

In the Standard Model of electroweak interactions and in the tree–approximation we calculate the rate and branching ratio of the neutron radiative β^-–decay with one–real photon emission by taking into account the contributions of the weak magnetism and proton recoil to order $1/m_p$ of the large proton mass m_p expansion. We find that the obtained contributions of the weak magnetism and proton recoil increase the rate and branching ratio of the neutron radiative β^-–decay by about 0.70 %. This is large compared with the contribution of the weak magnetism and proton recoil of about 0.16 % to the rate of the neutron β^-–decay, calculated in Phys. Rev. D **88**, 073002 (2013).

PACS numbers: 12.15.Ff, 13.15.+g, 23.40.Bw, 26.65.+t

arXiv:1701.04613v1 [hep-ph] 17 Jan 2017

I. INTRODUCTION

For a long time the radiative β^-–decay of a free neutron $n \to p + e^- + \bar{\nu}_e + \gamma$ was used in the analysis of the radiative corrections to the neutron β^-–decay for the cancellation of the infrared divergences, coming from the one–virtual photon exchanges [1]–[7]. For the first time to treat the radiative β^-–decay of a free neutron as a physical process was proposed by Gaponov and Khafizov [8], who made the first calculation of the energy spectrum and decay rate. Then, the radiative β^-–decay was reinvestigated in [9] and [10]. The branching ratio $\mathrm{BR}_{\beta\gamma} = 2.87 \times 10^{-3}$, calculated for the lifetime of the neutron $\tau_n = 879.6(1.1)\,\mathrm{s}$ in [10, 11] for the photon energy region $15\,\mathrm{keV} \leq \omega \leq 340\,\mathrm{keV}$, agrees well within one standard deviation with the first experimental values $\mathrm{BR}_{\beta\gamma} = 3.13(35) \times 10^{-3}$ and $\mathrm{BR}_{\beta\gamma} = 3.09(32) \times 10^{-3}$, measured by Nico *et al.* [12] and Cooper *et al.* [13], respectively. The experimental values agree also well with the result $\mathrm{BR}_{\beta\gamma} = 2.85 \times 10^{-3}$, calculated by Gardner [12] using the theoretical decay rate, published in [9]. The analysis of the T–odd momentum correlations even to order $O(\alpha^2)$ and the CP–violation by interactions beyond the Standard Model in the neutron radiative β^-–decay has been performed by Gardner and He in [14] and [15], respectively. Recently the new precise experimental values of the branching ratios of the radiative β^-–decay of a free neutron have been reported by the RDK II Collaboration Bales *et al.* [16]: $\mathrm{BR}_{\beta\gamma} = 0.00335 \pm 0.00005[\mathrm{stat}] \pm 0.00015[\mathrm{syst}]$ and $\mathrm{BR}_{\beta\gamma} = 0.00582 \pm 0.00023[\mathrm{stat}] \pm 0.00062[\mathrm{syst}]$, measured for the photon energy regions $14\,\mathrm{keV} \leq \omega \leq 782\,\mathrm{keV}$ and $0.4\,\mathrm{keV} \leq \omega \leq 14\,\mathrm{keV}$, respectively.

This paper is addressed to the calculation of the rate of the neutron radiative β^-–decay $n \to p + e^- + \bar{\nu}_e + \gamma$ in the Standard Model and in the tree–approximation by taking into account the contributions of the weak magnetism and proton recoil to order $1/M$, where $M = (m_n + m_p)/2$ is the averaged nucleon mass. The latter is important for a calculation of a robust theoretical background for the experimental analysis of interactions beyond the Standard Model [10, 11]. The paper is organized as follows. In section II we give the analytical expressions for the amplitude and rate of the neutron radiative β^--decay. In section III in Table I we adduce the results of the numerical analysis of the branching ratios of the neutron radiative β^-–decay for the experimental regions of photons energies. We discuss the obtained results and perspectives of the further theoretical analysis of the neutron radiative β^-–decay. In the Appendix we give the detailed calculation of the amplitude and rate of the neutron radiative β^-–decay in the tree–approximation.

*Electronic address: ivanov@kph.tuwien.ac.at
†Electronic address: roman.hoellwieser@gmail.com
‡Electronic address: natroitskaya@yandex.ru
§Electronic address: max.wellenzohn@gmail.com
¶Electronic address: berdnikov@spbstu.ru



FIG. 1: Feynman diagrams, defining the amplitude of the neutron radiative β^-–decay in the tree–approximation.

II. RADIATIVE β^-–DECAY OF NEUTRON IN THE TREE–APPROXIMATION TO NEXT–TO–LEADING ORDER IN THE LARGE M EXPANSION

In the Standard Model the neutron radiative β^-–decay is described by the following interactions

$$\mathcal{L}_{\rm int}(x) = \mathcal{L}_{\rm W}(x) + \mathcal{L}_{\rm em}(x). \tag{1}$$

Here $\mathcal{L}_{\rm W}(x)$ is the effective Lagrangian of low–energy $V-A$ interactions with a real axial coupling constant $\lambda = -1.2750(9)$ [17] and the contribution of the weak magnetism [10]

$$\mathcal{L}_{\rm W}(x) = -\frac{G_F}{\sqrt{2}}\,V_{ud}\,\Big\{[\bar{\psi}_p(x)\gamma_\mu(1+\lambda\gamma^5)\psi_n(x)] + \frac{\kappa}{2M}\partial^\nu[\bar{\psi}_p(x)\sigma_{\mu\nu}\psi_n(x)]\Big\}[\bar{\psi}_e(x)\gamma^\mu(1-\gamma^5)\psi_\nu(x)], \tag{2}$$

invariant under time reversal, where $G_F = 1.1664\times 10^{-11}\,{\rm MeV}^{-2}$ is the Fermi coupling constant, $|V_{ud}| = 0.97417(21)$ is the Cabibbo–Kobayashi–Maskawa matrix element and $\kappa = \kappa_p - \kappa_n = 3.7058$ is the isovector anomalous magnetic moment of the nucleon, defined by the anomalous magnetic moments of the proton $\kappa_p = 1.7928$ and the neutron $\kappa_n = -1.9130$ and measured in nuclear magneton [18]. Then, $\psi_p(x)$, $\psi_n(x)$, $\psi_e(x)$ and $\psi_\nu(x)$ are the field operators of the proton, neutron, electron and antineutrino, respectively, γ^μ, γ^5 and $\sigma^{\mu\nu} = \frac{i}{2}(\gamma^\mu\gamma^\nu - \gamma^\nu\gamma^\mu)$ are the Dirac matrices [19]. Then, $\mathcal{L}_{\rm em}(x)$ is the Lagrangian of the electromagnetic interaction

$$\mathcal{L}_{\rm em}(x) = -e\Big\{[\bar{\psi}_p(x)\gamma_\mu\psi_p(x)] - [\bar{\psi}_e(x)\gamma^\mu\psi_e(x)]\Big\}A_\mu(x), \tag{3}$$

where e is the proton electric charge, related to the fine–structure constant α by $e^2 = 4\pi\alpha$, and $A_\mu(x)$ is the electromagnetic potential [19].

In the tree–approximation the Feynman diagrams of the amplitude of the neutron radiative β^-–decay are shown in Fig. 1. The amplitude of the neutron radiative β^-–decay, defined by the diagrams in Fig. 1, we describe by the expression [10]

$$M(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'} = e\,\frac{G_F}{\sqrt{2}}\,V_{ud}\,\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}, \tag{4}$$

where $\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}$ is equal to

$$\begin{aligned}\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'} &= \Big[\bar{u}_p(\vec{k}_p,\sigma_p)\,\hat{\varepsilon}^*_{\lambda'}(k)\,\frac{1}{m_p - \hat{k}_p - \hat{k} - i0}\,O_\mu u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\gamma^\mu(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]\\ &- \Big[\bar{u}_p(\vec{k}_p,\sigma_p)O_\mu u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\hat{\varepsilon}^*_{\lambda'}(k)\,\frac{1}{m_e - \hat{k}_e - \hat{k} - i0}\,\gamma^\mu(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big].\end{aligned} \tag{5}$$

The matrix O_μ is given by [10]

$$O_\mu = \gamma_\mu(1+\lambda\gamma^5) + i\,\frac{\kappa}{2M}\,\sigma_{\mu\nu}(k_p - k_n)^\nu. \tag{6}$$

Then, $\bar{u}_p(\vec{k}_p,\sigma_p)$, $u_n(\vec{k}_n,\sigma_n)$, $\bar{u}_e(\vec{k}_e,\sigma_e)$ and $v_\nu(\vec{k}_\nu,+\frac{1}{2})$ are the Dirac wave functions of the free proton, neutron, electron and electron antineutrino with 3–momenta $\vec{k}_p$, $\vec{k}_n = \vec{0}$, $\vec{k}_e$ and $\vec{k}_\nu$ and polarizations $\sigma_p = \pm 1$, $\sigma_n = \pm 1$, $\sigma_e = \pm 1$ and $+\frac{1}{2}$ [10, 23], respectively, $\varepsilon^{*\alpha}_{\lambda'}(k)$ is the polarization vector of the photon in the polarization state $\lambda' = 1, 2$ with a 4–momentum k, obeying the constraint $\varepsilon^*_{\lambda'}(k)\cdot k = 0$. The amplitude Eq.(5) is gauge invariant. Indeed, one may show that, replacing $\varepsilon^{\alpha *}_{\lambda'}(k) \to k^\alpha$ and using the Dirac equations for the free proton and electron, the amplitude Eq.(5) vanishes.

The rate of the neutron radiative β^-–decay, described by the Feynman diagrams in Fig. 1 with photon from the energy region $\omega_{\rm min} \leq \omega \leq \omega_{\rm max}$, is equal to (see [10] and Eq.(A-33) of the Appendix)

$$\lambda_{\beta\gamma}(\omega_{\rm max},\omega_{\rm min}) = (1+3\lambda^2)\frac{\alpha}{\pi}\frac{G_F^2|V_{ud}|^2}{2\pi^3}\int_{\omega_{\rm min}}^{\omega_{\rm max}}\frac{d\omega}{\omega}\int_{m_e}^{E_0-\omega} dE_e\, E_e\sqrt{E_e^2-m_e^2}\,(E_0-E_e-\omega)^2\,F(E_e,Z=1)\,\rho_{\beta\gamma}(E_e,\omega), \tag{7}$$

where $E_0 = (m_n^2 - m_p^2 + m_e^2)/2m_n$ is the end–point energy of the electron energy–spectrum of the neutron β^-–decay [10] and $F(E_e, Z=1)$ is the relativistic Fermi function, describing Coulomb proton–electron final–state interaction. It is equal to [10]

$$F(E_e,Z=1) = \Big(1+\frac{1}{2}\gamma\Big)\,\frac{4(2r_p m_e\beta)^{2\gamma}}{\Gamma^2(3+2\gamma)}\,\frac{e^{\,\pi\alpha/\beta}}{(1-\beta^2)^{\gamma}}\,\Big|\Gamma\Big(1+\gamma+i\,\frac{\alpha}{\beta}\Big)\Big|^2, \tag{8}$$

where $\beta = k_e/E_e = \sqrt{E_e^2-m_e^2}/E_e$ is the electron velocity, $\gamma = \sqrt{1-\alpha^2}-1$, r_p is the electric radius of the proton and $\alpha = 1/137.036$ is the fine–structure constant. In numerical calculations we will use $r_p = 0.841\,{\rm fm}$ [22]. The function $\rho_{\beta\gamma}(E_e,\omega)$ is calculated in the Appendix. It is given by the integral

$$\begin{aligned}
\rho_{\beta\gamma}(E_e,\omega) &= \int\frac{d\Omega_{e\gamma}}{4\pi}\Big[1+2\,\frac{\omega}{M}\,\frac{E_e-\vec{k}_e\cdot\vec{n}}{E_0-E_e-\omega}+\frac{3}{M}\Big(E_e+\omega-\frac{1}{3}\,E_0\Big)+\frac{\lambda^2-2(\kappa+1)\lambda+1}{1+3\lambda^2}\,\frac{E_0-E_e-\omega}{M}\Big]\\
&\times\Big[\Big(1+\frac{\omega}{E_e}\Big)\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega^2}{E_e}\,\frac{1}{E_e-\vec{k}_e\cdot\vec{n}}\Big]+\frac{3\lambda^2-1}{1+3\lambda^2}\,\frac{1}{M}\,\Big\{\frac{k_e^2+\omega\vec{k}_e\cdot\vec{n}}{E_e}\,\Big(\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega}{E_e-\vec{k}_e\cdot\vec{n}}\Big)\\
&+(\omega+\vec{k}_e\cdot\vec{n}\,)\Big[\Big(1+\frac{\omega}{E_e}\Big)\frac{\omega}{E_e-\vec{k}_e\cdot\vec{n}}-\frac{m_e^2}{E_e}\,\frac{\omega}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}\Big]\Big\}-\frac{\lambda^2+2(\kappa+1)\lambda-1}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{k_e^2+\omega^2+2\omega\vec{k}_e\cdot\vec{n}}{E_e}\\
&\times\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega}{E_e}\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{E_e-\vec{k}_e\cdot\vec{n}}+\frac{\omega^2}{E_e}\,\frac{\omega+\vec{k}_e\cdot\vec{n}}{E_e-\vec{k}_e\cdot\vec{n}}\Big]-\frac{\lambda(\lambda-1)}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{\omega}{E_e}\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{E_e-\vec{k}_e\cdot\vec{n}}+3\,\frac{\omega^2}{E_e}\Big].
\end{aligned} \tag{9}$$

Here $d\Omega_{e\gamma}$ is an infinitesimal solid angle of the electron–photon momentum correlations $\vec{k}_e\cdot\vec{n} = k_e\cos\theta_{e\gamma}$, where $\vec{n} = \vec{k}/\omega$ is a unit vector along the photon 3–momentum [10, 11]. The results of the numerical analysis of the rate of the neutron radiative β^-–decay, calculated relative to the neutron lifetime $\tau_n = 879.6(1.1)\,{\rm s}$ [10, 11], we give and discuss in section III.

III. DISCUSSION AND CONCLUSION

Recent new experimental measurements of the branching ratio of the neutron radiative β^-–decay, reported by the RDK II Collaboration Bales *et al.* [16], have been the impetus for our theoretical analysis of the neutron radiative β^-–decay. We have performed the calculation of the amplitude of the neutron radiative β^-–decay in the tree–approximation to next–to–leading order in the large proton mass expansion. We have taken into account the complete set of contributions of the weak magnetism and proton recoil to order $1/M$, where M is the averaged nucleon mass. The obtained results we consider as a first step towards the precision theoretical analysis of the neutron radiative β^-–decay in the Standard Model to a relative order 10^{-3} [10, 11]. The detailed calculations of the amplitude and decay rate we give in the Appendix. The numerical values of the branching ratios of the neutron radiative β^-–decay, calculated relative to the neutron lifetime $\tau_n = 879.6(1.1)\,{\rm s}$ [10, 11], for the experimental regions of photon energies are adduced in the Table I.

ω [keV]	${\rm BR}_{\beta\gamma}$(Experiment)	${\rm BR}_{\beta\gamma}$(Theory)	${\rm BR}_{\beta\gamma}$ (Theory) M $\to\infty$	$\Delta{\rm BR}_{\beta\gamma}$ (Theory)
$15 \leq \omega \leq 340$	$(3.09 \pm 0.32)\times 10^{-3}$ [13]	2.89×10^{-3}	2.87×10^{-3}	0.70 %
$14 \leq \omega \leq 782$	$(3.35 \pm 0.05\,[{\rm stat}] \pm 0.15\,[{\rm syst}])\times 10^{-3}$ [16]	3.04×10^{-3}	3.02×10^{-3}	0.66 %
$0.4 \leq \omega \leq 14$	$(5.82 \pm 0.23\,[{\rm stat}] \pm 0.62\,[{\rm syst}])\times 10^{-3}$ [16]	5.08×10^{-3}	5.05×10^{-3}	0.60 %

TABLE I: Branching ratios of the radiative β^-–decay of the neutron for three photon energy regions, calculated for the lifetime of the neutron $\tau_n = 879.6(1.1)\,{\rm s}$ [10]. In the last column we give a relative contribution of the $1/M$ corrections, caused by the weak magnetism and proton recoil.

Of course, the numerical values of the branching ratios should not be practically changed if we would use the world averaged value of the lifetime of the neutron $\tau_n = 880.2(1.0)\,{\rm s}$ [18], which agrees perfectly well with the theoretical

one $\tau_n = 879.6(1.1)\,\mathrm{s}$ [10]. From the comparison of the branching ratios of the neutron radiative β^-–decay, calculated to leading order in the large proton mass expansion, one may see that the contributions of the weak magnetism and proton recoil make up of about 0.70 %, 0.66 % and 0.60 % for the photon energy regions $15\,\mathrm{keV} \leq \omega \leq 340\,\mathrm{keV}$, $14\,\mathrm{keV} \leq \omega \leq 782\,\mathrm{keV}$ and $0.4\,\mathrm{keV} \leq \omega \leq 14\,\mathrm{keV}$, respectively. Thus, at first glimpse, the contributions of the weak magnetism and proton recoil to the neutron radiative β^-–decays seem to be not very important. Moreover, that such contributions are small compared to the values of the statistic errors of the new experimental data 1.49 % and 3.95 % for the photon energy regions $14\,\mathrm{keV} \leq \omega \leq 782\,\mathrm{keV}$ and $0.4\,\mathrm{keV} \leq \omega \leq 14\,\mathrm{keV}$, respectively. The account for the systematic errors, making up of about 4.78 % and 11.34 % of the new experimental values, measured for the photon energy regions $14\,\mathrm{keV} \leq \omega \leq 782\,\mathrm{keV}$ and $0.4\,\mathrm{keV} \leq \omega \leq 14\,\mathrm{keV}$, respectively, makes the contribution of the weak magnetism and proton recoil fully intangible. Nevertheless, we would like to emphasize that the values of the $1/M$ corrections, caused by the weak magnetism and proton recoil, to the branching ratios of the neutron radiative β^-–decay are large compared by a factor of 4 to the contribution of the weak magnetism and proton recoil of about 0.16 % to the neutron lifetime, calculated to order $1/M$ in [10]. Thus, one may conclude that the weak magnetism and proton recoil, taken to order $1/M$, play more important role for the rate of the neutron radiative β^-–decay than for the rate of the neutron β^-–decay. On the whole they give a relative corrections of order 10^{-2}.

Combing the statistic and systematic errors the experimental values for the branching ratios we obtain $\mathrm{BR}^{(\mathrm{exp})}_{\beta\gamma} = 3.35(16)\times 10^{-3}$ and $\mathrm{BR}^{(\mathrm{exp})}_{\beta\gamma} = 5.82(66)\times 10^{-3}$ for the photon energy regions $14\,\mathrm{keV} \leq \omega \leq 782\,\mathrm{keV}$ and $0.4\,\mathrm{keV} \leq \omega \leq 14\,\mathrm{keV}$, respectively. One may see that the theoretical values of the branching ratios may agree with the experimental ones only within 2 and 1.2 standard deviations, respectively. The theoretical values of the branching ratios are below the experimental mean-values by about 9 % and 13 %. This leaves room for the analysis of other contributions to the neutron radiative β^-–decay. Following Bernard *et al.* [9] these contributions in the tree–approximation can be collected from the baryon resonances [18]. For example, the contribution of the $\Delta(1232)$–resonance with spin and parity $J^\pi = \frac{3}{2}^+$, the mass of which $m_\Delta \simeq 1232\,\mathrm{MeV}$ is not far from the proton mass [18], has been calculated by Bernard *et al.* [9]. According to Cooper *et al.* [13], the contribution of the $\Delta(1232)$–resonance makes up only of about 0.5 % of the branching ratio, measured for the photon energy region $15\,\mathrm{keV} \leq \omega \leq 340\,\mathrm{keV}$ [13]. This implies that other baryon resonances with heavier masses [18] should give contributions to the rate of the neutron radiative β^-–decay, which are small compared even to that by the $\Delta(1232)$–resonance.

Thus, one may expect that the theoretical analysis of the neutron radiative β^-–decay, which can be performed in the Standard Model and in the tree–approximation, may in principle change the rate of the neutron radiative β^-–decay not stronger than by about 1.5 % or even smaller. Hence, as a next step towards a precision theoretical analysis of the neutron radiative β^-–decay, which can be performed in the Standard Model, we may relate only to the analysis beyond the tree–approximation. To our point of view this should concern the radiative corrections to order $O(\alpha/\pi)$, calculated to leading order in the large proton mass expansion [10]. It is well–known [5] that the radiative corrections, calculated to order $O(\alpha/\pi)$ and to leading order in the large proton mass expansion, change the rate of the neutron β^-–decay by about 3.75 % [10]. Taking into account that the corrections of the weak magnetism and proton recoil of order $1/M$ to the neutron radiative β^-–decay rate are by a factor 4 large compared to the $1/M$ corrections to the rate of the neutron β^-–decay one may expect that the relative contribution of the radiative corrections of order $O(\alpha/\pi)$ to the rate of the neutron radiative β^-–decay can be also substantially enhanced. The first step in the direction of the account for the radiative corrections of order $O(\alpha/\pi)$ to the neutron radiative β^-–decay has been made by Gardner and He [14, 15]. However, the results, obtained by Gardner and He [14, 15], concern only the radiative corrections of order $O(\alpha/\pi)$ to T–odd momentum correlations in the neutron radiative β^-–decay. We are planning to give a detailed analysis of the radiative corrections of order $O(\alpha/\pi)$, allowing to describe the rate of the neutron radiative β^-–decay to order $O(\alpha^2/\pi^2)$, in our forthcoming publication.

IV. ACKNOWLEDGEMENTS

The work of A. N. Ivanov was supported by the Austrian "Fonds zur Förderung der Wissenschaftlichen Forschung" (FWF) under Contracts I689-N16, I862-N20 and P26781-N20 and by the ÖAW within the New Frontiers Groups Programme, NFP 2013/09. The work of R. Höllwieser was supported by the Erwin Schrödinger Fellowship program of the Austrian Science Fund FWF ("Fonds zur Förderung der wissenschaftlichen Forschung") under Contract No. J3425-N27. The work of M. Wellenzohn was supported by the MA 23 (FH-Call 16) under the project "Photonik - Stiftungsprofessur für Lehre".

Appendix A: Amplitude of radiative β^-–decay of neutron, described by Feynman diagrams in Fig. 1, with "weak magnetism" and proton recoil corrections to order $1/M$

The amplitude of the neutron radiative β^-–decay, given by Eq.(5), we transcribe into the form [10, 20]

$$\begin{aligned}\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'} &= \Big[\bar{u}_p(\vec{k}_p,\sigma_p)O_\mu u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\gamma^\mu(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big]\\ &-\Big[\bar{u}_p(\vec{k}_p,\sigma_p)\,Q_p\,\frac{1}{2k_p\cdot k}\,O_\mu u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\gamma^\mu(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big],\end{aligned}\tag{A-1}$$

where $\bar{u}_p(\vec{k}_p,\sigma_p)$, $u_n(\vec{k}_n,\sigma_n)$, $\bar{u}_e(\vec{k}_e,\sigma_e)$ and $v_\nu(\vec{k}_\nu,+\frac{1}{2})$ are the Dirac bispinor wave functions of the free proton, neutron, electron and electron antineutrino with 3–momenta $\vec{k}_p$, $\vec{k}_n=\vec{0}$, $\vec{k}_e$ and $\vec{k}_\nu$ and polarizations $\sigma_p=\pm1$, $\sigma_n=\pm1$, $\sigma_e=\pm1$ and $+\frac{1}{2}$ [10, 23], respectively, Q_e and Q_p are defined by

$$\begin{aligned}Q_e &= 2\varepsilon^*_{\lambda'}\cdot k_e+\hat{\varepsilon}^*_{\lambda'}\hat{k},\\ Q_p &= 2\varepsilon^*_{\lambda'}\cdot k_p+\hat{\varepsilon}^*_{\lambda'}\hat{k}.\end{aligned}\tag{A-2}$$

For the derivation of Eq.(A-1) we have used the Dirac equations for the free proton and electron. Replacing $\varepsilon^*_{\lambda'}\to k$ and using $k^2=0$ we get $\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)|_{\varepsilon^*_{\lambda'}\to k}=0$. The matrix Q_μ is given by Eq.(6). For the analysis of the neutron radiative β^-–decay in the non–relativistic approximation we have to rewrite the amplitude Eq.(A-1) in terms of time and space components of the matrix $O_\mu=(O_0,-\vec{O}\,)$ [10]. We get

$$\begin{aligned}\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'} &= \Big[\bar{u}_p(\vec{k}_p,\sigma_p)O_0u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\gamma^0(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big]\\ &-\Big[\bar{u}_p(\vec{k}_p,\sigma_p)\,\vec{O}\,u_n(\vec{k}_n,\sigma_n)\Big]\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big]\\ &-\Big[\bar{u}_p(\vec{k}_p,\sigma_p)Q_p\frac{1}{2k_p\cdot k}\,O_0u_n(\vec{k}_n,\sigma_n)\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\gamma^0(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big]\\ &+\Big[\bar{u}_p(\vec{k}_p,\sigma_p)\,\frac{1}{2k_p\cdot k}\,\vec{O}\,u_n(\vec{k}_n,\sigma_n)\Big]\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k}_\nu,+\frac{1}{2})\Big].\end{aligned}\tag{A-3}$$

The time O_0 and spatial $\vec{O}$ components of the matrix O_μ we determine to order $1/M$ in the large M expansion [10]

$$O_0=\begin{pmatrix}1 & \lambda+\dfrac{\kappa}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\\ -\lambda+\dfrac{\kappa}{2M}\,(\vec{\sigma}\cdot\vec{k}_p) & -1\end{pmatrix}\tag{A-4}$$

and

$$\vec{O}=\begin{pmatrix}\lambda\vec{\sigma}+i\,\dfrac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p) & \vec{\sigma}\,\Big(1-\dfrac{\kappa}{2M}\,(E_e+E_\nu+\omega)\Big)\\ -\,\vec{\sigma}\,\Big(1+\dfrac{\kappa}{2M}\,(E_e+E_\nu+\omega)\Big) & -\lambda\vec{\sigma}+i\dfrac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\end{pmatrix}.\tag{A-5}$$

For the derivation of Eq.(A-4) and Eq.(5) we have kept only the terms of order $1/M$ and used the energy conservation $m_n=E_p+E_e+E_\nu+\omega$, where m_n, $E_p=\sqrt{m_p^2+\vec{k}_p^{\,2}}$, E_e, E_ν and ω are the neutron mass and the proton, electron, antineutrino and photon energies, respectively. Then, $\vec{\sigma}$ are the Pauli 2×2 matrices [19].

For the calculation of the amplitude of the neutron radiative β^-–decay we define the Dirac bispinor wave functions of the neutron and the proton as follows

$$u_n(\vec{0},\sigma_n)=\sqrt{2m_n}\begin{pmatrix}\varphi_n\\ 0\end{pmatrix}\quad,\quad u_p(\vec{k}_p,\sigma_p)=\sqrt{E_p+m_p}\begin{pmatrix}\varphi_p\\ \dfrac{\vec{\sigma}\cdot\vec{k}_p}{E_p+m_p}\,\varphi_p\end{pmatrix},\tag{A-6}$$

where the Pauli spinorial wave functions φ_n and φ_p depend on the polarizations σ_n and σ_p, respectively. The matrix elements $[\bar{u}_p(\vec{k}_p,\sigma_p)O_0u_n(\vec{0},\sigma_n)]$ and $[\bar{u}_p(\vec{k}_p,\sigma_p)\vec{O}u_n(\vec{0},\sigma_n)]$ are equal to [10]

$$[\bar{u}_p(\vec{k}_p,\sigma_p)O_0u_n(\vec{0},\sigma_n)]=\sqrt{2m_n(E_p+m_p)}\,\Big\{[\varphi^\dagger_p\varphi_n]+\frac{\lambda}{2M}\,[\varphi^\dagger_p(\vec{\sigma}\cdot\vec{k}_p)\varphi_n]\Big\}\tag{A-7}$$

and

$$[\bar{u}_p(\vec{k}_p,\sigma_p)\vec{O}u_n(\vec{0},\sigma_n)] = \sqrt{2m_n(E_p+m_p)}\left\{\lambda[\varphi_p^\dagger\vec{\sigma}\,\varphi_n] + i\,\frac{\kappa}{2M}\,[\varphi_p^\dagger(\vec{\sigma}\times\vec{k}_p)\varphi_n] + \frac{1}{2M}\,[\varphi_p^\dagger(\vec{\sigma}\cdot\vec{k}_p)\vec{\sigma}\,\varphi_n]\right\}. \quad \text{(A-8)}$$

For the calculation of the matrix elements

$$[\bar{u}_p(\vec{k}_p,\sigma_p)Q_p\frac{1}{2k_p\cdot k}O_0u_n(\vec{0},\sigma_n)]\ ,\ [\bar{u}_p(\vec{k}_p,\sigma_p)Q_p\frac{1}{2k_p\cdot k}\vec{O}u_n(\vec{0},\sigma_n)] \quad \text{(A-9)}$$

we have to define the products $Q_p(1/2k_p\cdot k)\,Q_0$ and $Q_p(1/2k_p\cdot k)\,\vec{O}$. For this aim we define the matrix $Q_p(1/2k_p\cdot k)$ as follows

$$O_p\,\frac{1}{2k_p\cdot k} = \begin{pmatrix} \dfrac{2\varepsilon^*_{\lambda'}\cdot k_p}{2k_p\cdot k} + i\,\dfrac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2k_p\cdot k} & \dfrac{-\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)+\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2k_p\cdot k} \\ \dfrac{-\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)+\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2k_p\cdot k} & \dfrac{2\varepsilon^*_{\lambda'}\cdot k_p}{2k_p\cdot k} + i\,\dfrac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2k_p\cdot k} \end{pmatrix}, \quad \text{(A-10)}$$

where we have denoted $\varepsilon^{*\alpha}_{\lambda'} = (\varepsilon^{0*}_{\lambda'},\vec{\varepsilon}^{\,*}_{\lambda'})$ and $k^\alpha = (\omega,\vec{k}\,)$. Replacing $\varepsilon^{*\alpha}_{\lambda'} = (\varepsilon^{0*}_{\lambda'},\vec{\varepsilon}^{\,*}_{\lambda'}) \to k^\alpha = (\omega,\vec{k}\,)$ we get the matrix $Q_p(1/2k_p\cdot k)$ equal to a unit matrix $Q_p(1/2k_p\cdot k)|_{\varepsilon^{*\alpha}_{\lambda'}\to k} = 1$. Thus, the product $Q_p\,(1/2k_p\cdot k)\,Q_0$ is given by

$$O_p\frac{1}{2k_p\cdot k}Q_0 = \begin{pmatrix} \Big(O_p\dfrac{1}{2k_p\cdot k}Q_0\Big)_{11} & \Big(O_p\dfrac{1}{2k_p\cdot k}Q_0\Big)_{12} \\ \Big(O_p\dfrac{1}{2k_p\cdot k}Q_0\Big)_{21} & \Big(O_p\dfrac{1}{2k_p\cdot k}Q_0\Big)_{22} \end{pmatrix}, \quad \text{(A-11)}$$

where we have denoted

$$\begin{aligned}
\Big(O_p\frac{1}{2k_p\cdot k}Q_0\Big)_{11} &= \frac{\varepsilon^{0*}_{\lambda'}}{\omega} + i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega} + \lambda\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}, \\
\Big(O_p\frac{1}{2k_p\cdot k}Q_0\Big)_{12} &= \frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda+\frac{\kappa}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\Big) + i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega} + \frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}, \\
\Big(O_p\frac{1}{2k_p\cdot k}Q_0\Big)_{21} &= \frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(-\lambda+\frac{\kappa}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\Big) - i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega} - \frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}, \\
\Big(O_p\frac{1}{2k_p\cdot k}Q_0\Big)_{22} &= -\frac{\varepsilon^{0*}_{\lambda'}}{\omega} - i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega} - \lambda\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}.
\end{aligned} \quad \text{(A-12)}$$

In turn, the product $Q_p\,(1/2k_p\cdot k)\,\vec{O}$ reads

$$O_p\frac{1}{2k_p\cdot k}\vec{Q} = \begin{pmatrix} \Big(O_p\dfrac{1}{2k_p\cdot k}\vec{Q}\Big)_{11} & \Big(O_p\dfrac{1}{2k_p\cdot k}\vec{Q}\Big)_{12} \\ \Big(O_p\dfrac{1}{2k_p\cdot k}\vec{Q}\Big)_{21} & \Big(O_p\dfrac{1}{2k_p\cdot k}\vec{Q}\Big)_{22} \end{pmatrix}, \quad \text{(A-13)}$$

where we have denoted

$$\begin{aligned}
\Big(O_p\frac{1}{2k_p\cdot k}\vec{Q}\Big)_{11} &= \frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda\vec{\sigma} + i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\Big) + i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma} + \lambda\,\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}, \\
\Big(O_p\frac{1}{2k_p\cdot k}\vec{Q}\Big)_{12} &= \frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(1-\frac{\kappa}{2M}\,(E_e+E+\omega)\Big) + i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma} + \frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}, \\
\Big(O_p\frac{1}{2k_p\cdot k}\vec{Q}\Big)_{21} &= -\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(1+\frac{\kappa}{2M}\,(E_e+E+\omega)\Big) - i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma} - \lambda\,\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}, \\
\Big(O_p\frac{1}{2k_p\cdot k}\vec{Q}\Big)_{22} &= -\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda\vec{\sigma} + i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\Big) - i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma} - \frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}.
\end{aligned} \quad \text{(A-14)}$$

We would like to remind that because of the relation $\varepsilon^*_{\lambda'}\cdot k = 0$ the time–component of the polarization vector is equal to $\varepsilon^{0*}_{\lambda'} = (\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{k}\,)/\omega$. Replacing $\vec{\varepsilon}^{\,*}_{\lambda'} \to \vec{k}$ and using $|\vec{k}\,| = \omega$ we get $Q_p\,(1/2k_p\cdot k)\,Q_0|_{\vec{\varepsilon}^{\,*}_{\lambda'}\to\vec{k}} = Q_0$ and

$Q_p\,(1/2k_p\cdot k)\,\vec{O}\,|_{\vec{\varepsilon}^{\,*}_{\lambda'}\to\vec{k}}=\vec{O}$, respectively. Now we may define the matrix elements

$$[\bar{u}_p(\vec{k}_p,\sigma_p)Q_p\frac{1}{2k_p\cdot k}O_0u_n(\vec{0},\sigma_n)]=\sqrt{2m_n(E_p+m_p)}\,\Big\{\varphi_p^\dagger\Big(O_p\frac{1}{2k_p\cdot k}O_0\Big)_{11}\varphi_n+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,\varphi_p^\dagger(\vec{\sigma}\cdot\vec{k}_p)\varphi_n\Big\}=$$
$$=\sqrt{2m_n(E_p+m_p)}\,\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}+i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}+\lambda\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_n\Big] \tag{A-15}$$

and

$$[\bar{u}_p(\vec{k}_p,\sigma_p)Q_p\frac{1}{2k_p\cdot k}\vec{O}u_n(\vec{0},\sigma_n)]=\sqrt{2m_n(E_p+m_p)}\,\Big\{\varphi_p^\dagger\Big(O_p\frac{1}{2k_p\cdot k}\vec{Q}\Big)_{11}\varphi_n+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,\varphi_p^\dagger(\vec{\sigma}\cdot\vec{k}_p)\varphi_n\Big\}=$$
$$=\sqrt{2m_n(E_p+m_p)}\,\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda\vec{\sigma}+i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\Big)+i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}$$
$$+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\,\vec{\sigma}\Big)\varphi_n\Big]. \tag{A-16}$$

The amplitude of the neutron radiative β^-–decay, calculated to order $1/M$, caused by the weak magnetism and proton recoil, is equal to

$$\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}=\sqrt{2m_n(E_p+m_p)}\,\Big\{\Big[\varphi_p^\dagger\Big(1+\frac{\lambda}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_n\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\gamma^0(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]$$
$$-\Big[\varphi_p^\dagger\Big(\lambda\,\vec{\sigma}+i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)+\frac{1}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\vec{\sigma}\Big)\varphi_n\Big]\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]$$
$$-\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}+i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}+\lambda\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_n\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\gamma^0(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]$$
$$+\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda\vec{\sigma}+i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\Big)+i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\,\vec{\sigma}\Big)\varphi_n\Big]$$
$$\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]\Big\}. \tag{A-17}$$

Since the factor $\sqrt{2m_m(E_p+m_p)}$ to order $1/M$ is equal to

$$\sqrt{2m_m(E_p+m_p)}=2m_n\Big(1-\frac{E_e+E_\nu+\omega}{2M}\Big), \tag{A-18}$$

the amplitude Eq.(A-17) can be transcribed into the form

$$\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}=2m_n\,\Big\{\Big[\varphi_p^\dagger\Big(1-\frac{E_e+E_\nu+\omega}{2M}+\frac{\lambda}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_n\Big]\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\gamma^0(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]$$
$$-\Big[\varphi_p^\dagger\Big(\lambda\,\vec{\sigma}-\lambda\,\frac{E_e+E_\nu+\omega}{2M}\,\vec{\sigma}+i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)+\frac{1}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\vec{\sigma}\Big)\varphi_n\Big]\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\frac{1}{2k_e\cdot k}\,Q_e\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]$$
$$-\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}-\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\,\frac{E_e+E_\nu+\omega}{2M}+i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}+\lambda\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_n\Big]$$
$$\times\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\gamma^0(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]+\Big[\varphi_p^\dagger\Big(\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(\lambda\,\vec{\sigma}-\lambda\,\frac{E_e+E_\nu+\omega}{2M}\,\vec{\sigma}+i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)\Big)+i\,\lambda\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)}{2M\omega}\,\vec{\sigma}$$
$$+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}\,\vec{\sigma}+\lambda\,\frac{\varepsilon^{0*}_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\,\vec{\sigma}\Big)\varphi_n\Big]\cdot\Big[\bar{u}_e(\vec{k}_e,\sigma_e)\,\vec{\gamma}\,(1-\gamma^5)v_\nu(\vec{k},+\frac{1}{2})\Big]\Big\}. \tag{A-19}$$

The hermitian conjugate amplitude is equal to

$$\mathcal{M}^\dagger(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}=$$
$$=2m_n\,\Big\{\Big[\varphi_n^\dagger\Big(1-\frac{E_e+E_\nu+\omega}{2M}+\frac{\lambda}{2M}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_p\Big]\Big[\bar{v}_\nu(\vec{k},+\frac{1}{2})\,\frac{1}{2k_e\cdot k}\,\gamma^0\,\bar{Q}_e(1-\gamma^5)u_e(\vec{k}_e,\sigma_e)\Big]$$
$$-\Big[\varphi_n^\dagger\Big(\lambda\,\vec{\sigma}-\lambda\,\frac{E_e+E_\nu+\omega}{2M}\,\vec{\sigma}-i\,\frac{\kappa}{2M}\,(\vec{\sigma}\times\vec{k}_p)+\frac{1}{2M}\,\vec{\sigma}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_p\Big]\cdot\Big[\bar{v}_\nu(\vec{k},+\frac{1}{2})\,\frac{1}{2k_e\cdot k}\,\vec{\gamma}\,\bar{Q}_e\,(1-\gamma^5)u_e(\vec{k}_e,\sigma_e)\Big]$$
$$-\Big[\varphi_n^\dagger\Big(\frac{\varepsilon^0_{\lambda'}}{\omega}-\frac{\varepsilon^0_{\lambda'}}{\omega}\,\frac{E_e+E_\nu+\omega}{2M}-i\,\frac{\vec{\sigma}\cdot(\vec{\varepsilon}_{\lambda'}\times\vec{k}\,)}{2M\omega}+\lambda\frac{\varepsilon^0_{\lambda'}(\vec{\sigma}\cdot\vec{k}\,)-\omega(\vec{\varepsilon}_{\lambda'}\cdot\vec{\sigma}\,)}{2M\omega}+\lambda\,\frac{\varepsilon^0_{\lambda'}}{2M\omega}\,(\vec{\sigma}\cdot\vec{k}_p)\Big)\varphi_p\Big]$$

$$\times\Big[\bar v_\nu(\vec k,+\frac{1}{2})\,\gamma^0(1-\gamma^5)u_e(\vec k_e,\sigma_e)\Big]+\Big[\varphi_n^\dagger\Big(\frac{\varepsilon^0_{\lambda'}}{\omega}\Big(\lambda\,\vec\sigma-\lambda\,\frac{E_e+E_\nu+\omega}{2M}\,\vec\sigma-i\,\frac{\kappa}{2M}\,(\vec\sigma\times\vec k_p)\Big)-i\,\lambda\,\vec\sigma\,\frac{\vec\sigma\cdot(\vec\varepsilon_{\lambda'}\times\vec k\,)}{2M\omega}$$
$$+\lambda\,\vec\sigma\,\frac{\varepsilon^0_{\lambda'}(\vec\sigma\cdot\vec k\,)-\omega(\vec\varepsilon_{\lambda'}\cdot\vec\sigma\,)}{2M\omega}+\lambda\,\frac{\varepsilon^0_{\lambda'}}{2M\omega}\,\vec\sigma\,(\vec\sigma\cdot\vec k_p)\Big)\varphi_p\Big]\cdot\Big[\bar v_\nu(\vec k,+\frac{1}{2})\,\vec\gamma\,(1-\gamma^5)u_e(\vec k_e,\sigma_e)\Big]\Big\},\qquad\text{(A-20)}$$

where $\bar Q_e=2k_e\cdot k+\hat k\hat\varepsilon_{\lambda'}$ [10]. The squared absolute value of the amplitude Eq.(A-19) summed over the polarizations of the proton and electron and averaged over polarizations of the neutron is equal to

$$\sum_{\rm pol}\frac{|\mathcal{M}(n\to pe^-\bar\nu_e\gamma)_{\lambda'}|^2}{4m_n^2}=\frac{1}{(k_e\cdot k)^2}\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,{\rm tr}\{(\hat k_e+m_e)\,Q_e\gamma^0\hat k_\nu\gamma^0\bar Q_e(1-\gamma^5)\}-\frac{1}{(k_e\cdot k)^2}\,\frac{\lambda^2+1}{2M}$$
$$\times\vec k_p\cdot{\rm tr}\{(\hat k_e+m_e)\,Q_e\gamma^0\hat k_\nu\vec\gamma\,\bar Q_e(1-\gamma^5)\}-2\,\frac{\varepsilon^0_{\lambda'}}{\omega}\,\frac{1}{k_e\cdot k}\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,{\rm tr}\{(\hat k_e+m_e)\,Q_e\gamma^0\hat k_\nu\gamma^0\,(1-\gamma^5)\}$$
$$+\frac{1}{k_e\cdot k}\Big(\frac{\varepsilon^0_{\lambda'}}{\omega}\,\frac{\lambda^2+\lambda}{M}\,\vec k_p+\frac{\lambda}{M\omega}\,(\varepsilon^0_{\lambda'}\vec k-\omega\vec\varepsilon_{\lambda'})-\frac{\lambda}{M\omega}\,i\,(\vec\varepsilon_{\lambda'}\times\vec k\,)\Big)\cdot{\rm tr}\{(\hat k_e+m_e)\,Q_e\gamma^0\hat k_\nu\vec\gamma\,(1-\gamma^5)\}-\frac{1}{(k_e\cdot k)^2}\,\frac{\lambda^2+1}{2M}$$
$$\times\vec k_p\cdot{\rm tr}\{(\hat k_e+m_e)\,Q_e\vec\gamma\,\hat k_\nu\gamma^0\bar Q_e(1-\gamma^5)\}+\frac{1}{(k_e\cdot k)^2}\,\Big[\lambda^2\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\delta^{ij}+\frac{\lambda\,(\kappa+1)}{M}\,i\,\varepsilon^{ij\ell}(\vec k_p)^\ell\Big]$$
$$\times{\rm tr}\{(\hat k_e+m_e)\,Q_e\vec\gamma^{\,i}\,\hat k_\nu\vec\gamma^{\,j}\,\bar Q_e(1-\gamma^5)\}+\frac{1}{k_e\cdot k}\Big(\frac{\lambda^2+1}{M\omega}\,\varepsilon^0_{\lambda'}\vec k_p+\frac{\lambda^2}{M\omega}\,(\varepsilon^0_{\lambda'}\vec k-\omega\vec\varepsilon_{\lambda'})-\frac{\lambda}{M\omega}\,i\,(\vec\varepsilon_{\lambda'}\times\vec k\,)\Big)$$
$$\cdot{\rm tr}\{(\hat k_e+m_e)\,Q_e\vec\gamma\,\hat k_\nu\gamma^0(1-\gamma^5)\}-2\,\frac{1}{k_e\cdot k}\,\Big[\lambda^2\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\frac{\varepsilon^0_{\lambda'}}{\omega}\,\delta^{ij}+\frac{\lambda^2}{2M\omega}\Big(\vec\varepsilon^{\,i}_{\lambda'}\vec k^{\,j}-\vec\varepsilon^{\,j}_{\lambda'}\vec k^{\,i}\Big)$$
$$+\frac{\lambda^2+(2\kappa+1)\lambda}{2M\omega}\,\varepsilon^0_{\lambda'}\,i\,\varepsilon^{ij\ell}(\vec k_p)^\ell+\frac{\lambda^2}{2M\omega}\,i\,\varepsilon^{ij\ell}\Big(\varepsilon^0_{\lambda'}\vec k^{\,\ell}-\omega\vec\varepsilon^{\,\ell}_{\lambda'}\Big)\Big]{\rm tr}\{(\hat k_e+m_e)\,Q_e\vec\gamma^{\,i}\,\hat k_\nu\vec\gamma^{\,j}\,(1-\gamma^5)\}$$
$$-2\,\frac{1}{k_e\cdot k}\,\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,{\rm tr}\{(\hat k_e+m_e)\,\gamma^0\hat k_\nu\gamma^0\bar Q_e\,(1-\gamma^5)\}+\frac{1}{k_e\cdot k}\,\Big(\frac{\lambda^2+1}{M\omega}\,\varepsilon^{0*}_{\lambda'}\vec k_p+\frac{\lambda^2}{M\omega}\,(\varepsilon^{0*}_{\lambda'}\vec k-\omega\vec\varepsilon^{\,*}_{\lambda'})$$
$$+i\,\frac{\lambda}{M\omega}\,(\vec\varepsilon^{\,*}_{\lambda'}\times\vec k\,)\Big)\cdot{\rm tr}\{(\hat k_e+m_e)\,\gamma^0\,\hat k_\nu\vec\gamma\,\bar Q_e(1-\gamma^5)\}+4\,\frac{\varepsilon^{0*}_{\lambda'}\varepsilon^0_{\lambda'}}{\omega^2}\,\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,{\rm tr}\{(\hat k_e+m_e)\,\gamma^0\hat k_\nu\gamma^0(1-\gamma^5)\}$$
$$-2\,\Big[\frac{\lambda^2+\lambda}{M\omega^2}\,\varepsilon^{0*}_{\lambda'}\varepsilon^0_{\lambda'}\vec k_p+\frac{\lambda^2}{M\omega^2}\,(\varepsilon^{0*}_{\lambda'}\vec k-\omega\vec\varepsilon^{\,*}_{\lambda'})\varepsilon^0_{\lambda'}+\frac{\lambda}{M\omega^2}\,\varepsilon^{0*}_{\lambda'}(\varepsilon^0_{\lambda'}\vec k-\omega\vec\varepsilon_{\lambda'})+\frac{\lambda}{M\omega^2}\,i\,\Big((\vec\varepsilon^{\,*}_{\lambda'}\times\vec k\,)\varepsilon^0_{\lambda'}-\varepsilon^{0*}_{\lambda'}(\vec\varepsilon_{\lambda'}\times\vec k\,)\Big)\Big]$$
$$\cdot{\rm tr}\{(\hat k_e+m_e)\,\gamma^0\,\hat k_\nu\vec\gamma\,(1-\gamma^5)\}+\frac{1}{k_e\cdot k}\,\Big(\frac{\lambda^2+\lambda}{M\omega}\,\varepsilon^{0*}_{\lambda'}\,\vec k_p+\frac{\lambda}{M\omega}\,(\varepsilon^{0*}_{\lambda'}\,\vec k-\omega\,\vec\varepsilon^{\,*}_{\lambda'})+\frac{\lambda}{M\omega}\,i\,(\vec\varepsilon^{\,*}_{\lambda'}\times\vec k\,)\Big)$$
$$\cdot{\rm tr}\{(\hat k_e+m_e)\,\vec\gamma\,\hat k_\nu\gamma^0\,\bar Q_e(1-\gamma^5)\}-2\,\frac{1}{k_e\cdot k}\,\Big[\lambda^2\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\frac{\varepsilon^{0*}_{\lambda'}}{\omega}\,\delta^{ij}-\frac{\lambda^2}{2M\omega}\Big(\vec\varepsilon^{\,i*}_{\lambda'}\vec k^{\,j}-\vec\varepsilon^{\,j*}_{\lambda'}\vec k^{\,i}\Big)$$
$$+\frac{\lambda^2+(2\kappa+1)\lambda}{2M\omega}\,\varepsilon^{0*}_{\lambda'}\,i\,\varepsilon^{ij\ell}\vec k^{\,\ell}_p+\frac{\lambda^2}{2M\omega}\,i\,\varepsilon^{ij\ell}\Big(\varepsilon^{0*}_{\lambda'}\vec k^{\,\ell}-\omega\vec\varepsilon^{\,\ell*}_{\lambda'}\Big)\Big]{\rm tr}\{(\hat k_e+m_e)\,\vec\gamma^{\,i}\,\hat k_\nu\vec\gamma^{\,j}\,\bar Q_e(1-\gamma^5)\}$$
$$-2\,\Big[\frac{\lambda^2+\lambda}{M\omega^2}\,\varepsilon^{0*}_{\lambda'}\varepsilon^0_{\lambda'}\vec k_p+\frac{\lambda^2}{M\omega^2}\,\varepsilon^{0*}_{\lambda'}(\varepsilon^0_{\lambda'}\vec k-\omega\vec\varepsilon_{\lambda'})+\frac{\lambda}{M\omega^2}\,(\varepsilon^{0*}_{\lambda'}\vec k-\omega\vec\varepsilon^{\,*}_{\lambda'})\varepsilon^0_{\lambda'}-\frac{\lambda}{M\omega^2}\,i\,\Big(\varepsilon^{0*}_{\lambda'}(\vec\varepsilon_{\lambda'}\times\vec k\,)-(\vec\varepsilon^{\,*}_{\lambda'}\times\vec k\,)\varepsilon^0_{\lambda'}\Big)\Big]$$
$$\cdot{\rm tr}\{(\hat k_e+m_e)\,\vec\gamma\,\hat k_\nu\gamma^0(1-\gamma^5)\}+4\,\Big\{\lambda^2\frac{\varepsilon^{0*}_{\lambda'}\varepsilon^0_{\lambda'}}{\omega^2}\,\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\delta^{ij}+i\,\frac{\lambda^2+\kappa\lambda}{M\omega^2}\,\varepsilon^{0*}_{\lambda'}\varepsilon^0_{\lambda'}\,\varepsilon^{ij\ell}(\vec k_p)^\ell$$
$$-\frac{\lambda^2}{2M\omega^2}\,\Big[\Big(\vec\varepsilon^{\,i*}_{\lambda'}\vec k^{\,j}-\vec\varepsilon^{\,j*}_{\lambda'}\vec k^{\,i}\Big)\varepsilon^0_{\lambda'}-\varepsilon^{0*}_{\lambda'}\Big(\vec\varepsilon^{\,i}_{\lambda'}\vec k^{\,j}-\vec\varepsilon^{\,j}_{\lambda'}\vec k^{\,i}\Big)\Big]$$
$$+\frac{\lambda^2}{2M\omega^2}\,i\,\varepsilon^{ij\ell}\Big[\Big(\varepsilon^{0*}_{\lambda'}\vec k^{\,\ell}-\omega\vec\varepsilon^{\,\ell*}_{\lambda'}\Big)\varepsilon^0_{\lambda'}+\varepsilon^{0*}_{\lambda'}\Big(\varepsilon^0_{\lambda'}\vec k^{\,\ell}-\omega\vec\varepsilon^{\,\ell}_{\lambda'}\Big)\Big]\Big\}\,{\rm tr}\{(\hat k_e+m_e)\vec\gamma^{\,i}\hat k_\nu\vec\gamma^{\,j}(1-\gamma^5)\}.\qquad\text{(A-21)}$$

One may show that the expression in Eq.(A-21) is gauge invariant. For this aim one has to replace $\varepsilon^*_{\lambda'}\to k$ and $\varepsilon_{\lambda'}\to k$ and collect the coefficients in font of the traces ${\rm tr}\{(\hat k_e+m_e)\gamma^\alpha\hat k_\nu\gamma^\beta(1-\gamma^5)\}$. Now we may proceed to the calculation of the traces over electron and antineutrino degrees of freedom. Using the following relations [10] (see Appendix B of Ref.[10])

$$\gamma^\alpha\gamma^\nu\gamma^\mu=\gamma^\alpha\eta^{\nu\mu}-\gamma^\nu\eta^{\mu\alpha}+\gamma^\mu\eta^{\alpha\nu}+i\,\varepsilon^{\alpha\nu\mu\beta}\,\gamma_\beta\gamma^5,\qquad\text{(A-22)}$$

where $\eta^{\alpha\beta}$ is the Minkowski metric tensor and $\varepsilon^{\alpha\nu\mu\beta}$ is the Levi–Civita tensor defined by $\varepsilon^{0123}=1$ and $\varepsilon_{\alpha\nu\mu\beta}=-\varepsilon^{\alpha\nu\mu\beta}$ [19] and [10] (see Appendix B of Ref.[10]), we obtain

$$\frac{1}{4}\,{\rm tr}\{\hat k_eQ_e\gamma^\mu\bar Q_e(1-\gamma^5)\}=4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(k_e+k)^\mu-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,k^\mu$$

$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^{\mu}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^{\mu *}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{\mu\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,k^\mu\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu} \qquad \text{(A-23)}$$

for a photon on–mass shell $k^2=0$. Then, we get

$$\frac{1}{4}\,\mathrm{tr}\{\hat{k}_e Q_e\gamma^\mu(1-\gamma^5)\}=(\varepsilon^*_{\lambda'}\cdot k_e)(2k_e+k)^\mu-(k_e\cdot k)\varepsilon^{\mu *}_{\lambda'}-i\,\varepsilon^{\mu\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu},$$
$$\frac{1}{4}\,\mathrm{tr}\{\hat{k}_e\gamma^\mu\bar{Q}_e(1-\gamma^5)\}=(\varepsilon_{\lambda'}\cdot k_e)(2k_e+k)^\mu-(k_e\cdot k)\varepsilon^{\mu}_{\lambda'}+i\,\varepsilon^{\mu\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}. \qquad \text{(A-24)}$$

Using Eq.(A-23) and Eq.(A-24) for the traces over the electron and antineutrino degrees of freedom, containing both Q_e and $\bar{Q}_e$ matrices, we obtain the following expressions

$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(E_e+\omega)-2\,\omega\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{0\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\omega\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+\vec{k}^{\,i}_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^i-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,i}$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,(\vec{\varepsilon}^{\,i}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i*}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{i\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\vec{k}^{\,i}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\vec{\gamma}^{\,i}\,\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^i-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,i}$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i*}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{i\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\vec{k}^{\,i}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+\vec{k}^{\,i}_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(E_e+\omega)-2\,\omega\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{0\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\omega\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+i\,\varepsilon^{ij\ell}\vec{k}^{\,j}_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^\ell-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,(\vec{k}\,)^\ell$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,(\vec{\varepsilon}^{\,\ell}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell*}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{\ell\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\vec{k}^{\,\ell}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\,\hat{k}_\nu\gamma^0\,\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^i-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,i}$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,(\vec{\varepsilon}^{\,i}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i*}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{i\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\vec{k}^{\,i}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+\vec{k}^{\,i}_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(E_e+\omega)-2\,\omega\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{0\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\omega\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]-i\,\varepsilon^{ij\ell}\vec{k}^{\,j}_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^\ell-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,\ell}$$
$$-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell*}_{\lambda'}\right)(k_e\cdot k)+2\,i\,\varepsilon^{\ell\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}$$
$$+2\,i\,\vec{k}^{\,\ell}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}=$$
$$=E_\nu\frac{1}{4}\,\mathrm{tr}\{\hat{k}_e\,Q_e\vec{\gamma}^{\,i}\,\gamma^0\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}-\vec{k}^{\,\ell}_\nu\frac{1}{4}\,\mathrm{tr}\{\hat{k}_e\,Q_e\vec{\gamma}^{\,i}\,\vec{\gamma}^{\,\ell}\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}=$$
$$=\delta^{ij}\,E_\nu\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(E_e+\omega)-2\,\omega\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)-2\left((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}\right)(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{0\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,\omega\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]-i\,\varepsilon^{ij\ell}E_\nu$$

$$\times\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^\ell-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,\ell}-2\,\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell*}_{\lambda'}\Big)\,(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{\ell\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,(\vec{k}\,)^\ell\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]-\delta^{ij}\,\vec{k}^{\,\ell}_\nu$$
$$\times\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^\ell-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,\ell}-2\,\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,\ell*}_{\lambda'}\Big)\,(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{\ell\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,\vec{k}^{\,\ell}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+\vec{k}^{\,i}_\nu$$
$$\times\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^j-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,j}-2\,\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,j}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,j*}_{\lambda'}\Big)\,(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{j\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,\vec{k}^{\,j}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]+\vec{k}^{\,j}_\nu$$
$$\times\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(\vec{k}_e+\vec{k}\,)^i-2\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)\,\vec{k}^{\,i}-2\,\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i}_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\vec{\varepsilon}^{\,i*}_{\lambda'}\Big)\,(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{i\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,\vec{k}^{\,i}\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big]-i\,\varepsilon^{ij\ell}\vec{k}^{\,\ell}_\nu$$
$$\times\Big[4(\varepsilon^*_{\lambda'}\cdot k_e)(\varepsilon_{\lambda'}\cdot k_e)\,(E_e+\omega)-2\,\omega\,(\varepsilon^*_{\lambda'}\cdot\varepsilon_{\lambda'})\,(k_e\cdot k)-2\,\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}+(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^0_{\lambda'}\Big)\,(k_e\cdot k)$$
$$+2\,i\,\varepsilon^{0\alpha\beta\nu}\Big((\varepsilon^*_{\lambda'}\cdot k_e)\,\varepsilon_{\lambda'\alpha}-(\varepsilon_{\lambda'}\cdot k_e)\,\varepsilon^*_{\lambda'\alpha}\Big)\,k_\beta k_{e\nu}+2\,i\,\omega\,\varepsilon^{\alpha\beta\rho\nu}\,\varepsilon^*_{\lambda'\alpha}\,\varepsilon_{\lambda'\beta}\,k_\rho\,k_{e\nu}\Big] \tag{A-25}$$

For the traces over the electron and antineutrino degrees of freedom, containing either Q_e or $\bar{Q}_e$, we get

$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\gamma^0(1-\gamma^5)\}=E_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^{0*}_{\lambda'}-i\,\varepsilon^{0\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i*}_{\lambda'}-i\,\varepsilon^{i\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\vec{\gamma}^{\,i}(1-\gamma^5)\}=E_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i*}_{\lambda'}-i\,\varepsilon^{i\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^{0*}_{\lambda'}-i\,\varepsilon^{0\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+i\,\varepsilon^{ij\ell}\vec{k}^{\,j}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell*}_{\lambda'}-i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\hat{k}_\nu\gamma^0(1-\gamma^5)\}=E_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i*}_{\lambda'}-i\,\varepsilon^{i\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^{0*}_{\lambda'}-i\,\varepsilon^{0\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$-i\,\varepsilon^{ij\ell}\vec{k}^{\,j}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell*}_{\lambda'}-i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\hat{k}_\nu\vec{\gamma}^{\,j}(1-\gamma^5)\}=\delta^{ij}\,E_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^{0*}_{\lambda'}-i\,\varepsilon^{0\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$-i\,\varepsilon^{ij\ell}\,E_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell*}_{\lambda'}-i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$-\delta^{ij}\,\vec{k}^{\,\ell}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell*}_{\lambda'}-i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^j-(k_e\cdot k)\vec{\varepsilon}^{\,j*}_{\lambda'}-i\,\varepsilon^{j\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,j}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i*}_{\lambda'}-i\,\varepsilon^{i\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$-i\,\varepsilon^{ij\ell}\,\vec{k}^{\,\ell}_\nu\Big[(\varepsilon^*_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^{0*}_{\lambda'}-i\,\varepsilon^{0\alpha\beta\nu}\varepsilon^*_{\lambda'\alpha}k_\beta k_{e\nu}\Big] \tag{A-26}$$

and

$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^0_{\lambda'}+i\,\varepsilon^{0\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i}_{\lambda'}+i\,\varepsilon^{i\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big],$$
$$\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\hat{k}_\nu\vec{\gamma}^{\,i}\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k}\,)^i-(k_e\cdot k)\vec{\varepsilon}^{\,i}_{\lambda'}+i\,\varepsilon^{i\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$
$$+\vec{k}^{\,i}_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^0_{\lambda'}+i\,\varepsilon^{0\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]$$

$$
\begin{aligned}
&-i\,\varepsilon^{ij\ell}\vec{k}_\nu^{\,j}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell}_{\lambda'}+i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big],\\
&\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^i-(k_e\cdot k)\vec{\varepsilon}^{\,i}_{\lambda'}+i\,\varepsilon^{i\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&+\vec{k}_\nu^{\,i}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^0_{\lambda'}+i\,\varepsilon^{0\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&-i\,\varepsilon^{ij\ell}\vec{k}_\nu^{\,j}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell}_{\lambda'}+i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big],\\
&\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\hat{k}_\nu\vec{\gamma}^{\,j}\bar{Q}_e(1-\gamma^5)\}=\delta^{ij}\,E_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^0_{\lambda'}+i\,\varepsilon^{0\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&-i\,\varepsilon^{ij\ell}\,E_\nu\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell}_{\lambda'}+i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&-\delta^{ij}\,\vec{k}_\nu^{\,\ell}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^\ell-(k_e\cdot k)\vec{\varepsilon}^{\,\ell}_{\lambda'}+i\,\varepsilon^{\ell\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&+\vec{k}_\nu^{\,i}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^j-(k_e\cdot k)\varepsilon^j_{\lambda'}+i\,\varepsilon^{j\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&+\vec{k}_\nu^{\,j}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2\vec{k}_e+\vec{k})^i-(k_e\cdot k)\vec{\varepsilon}^{\,i}_{\lambda'}+i\,\varepsilon^{i\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]\\
&-i\,\varepsilon^{ij\ell}\,\vec{k}_\nu^{\,\ell}\Big[(\varepsilon_{\lambda'}\cdot k_e)(2E_e+\omega)-(k_e\cdot k)\varepsilon^0_{\lambda'}+i\,\varepsilon^{0\alpha\beta\nu}\varepsilon_{\lambda'\alpha}k_\beta k_{e\nu}\Big]. \qquad \text{(A-27)}
\end{aligned}
$$

In turn, for the traces without Q_e and $\bar{Q}_e$ we obtain the following expressions

$$
\begin{aligned}
\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\hat{k}_\nu\gamma^0(1-\gamma^5)\} &= 3E_eE_\nu+\vec{k}_e\cdot\vec{k}_\nu,\\
\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\hat{k}_\nu\vec{\gamma}^{\,i}(1-\gamma^5)\} &= E_e(\vec{k}_\nu)^i+E_\nu(\vec{k}_e)^i-i\,\varepsilon^{ij\ell}(\vec{k}_e)^j(\vec{k}_\nu)^\ell,\\
\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\hat{k}_\nu\gamma^0(1-\gamma^5)\} &= E_e(\vec{k}_\nu)^i+E_\nu(\vec{k}_e)^i+i\,\varepsilon^{ij\ell}(\vec{k}_e)^j(\vec{k}_\nu)^\ell,\\
\frac{1}{4}\,\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\hat{k}_\nu\vec{\gamma}^{\,j}(1-\gamma^5)\} &= (\vec{k}_e)^i(\vec{k}_\nu)^j+(\vec{k}_e)^j(\vec{k}_\nu)^i+\delta^{ij}(E_eE_\nu-\vec{k}_e\cdot\vec{k}_\nu)\\
&\quad+\ i\,\varepsilon^{ij\ell}(E_e(\vec{k}_\nu)^\ell-E_\nu(\vec{k}_e)^\ell). \qquad \text{(A-28)}
\end{aligned}
$$

Now we may sum over the photon polarizations. Because of gauge invariance of Eq.(A-21) one may use any gauge. However, it is obvious that one has to sum over only physical degrees of freedom of real photons, which are defined by the polarization vector $\varepsilon^*_{\lambda'}=(0,\vec{\varepsilon}^{\,*}_{\lambda'})$ [21] (see also [20]). The polarization vector $\vec{\varepsilon}^{\,*}_{\lambda'}$ has the following properties [21] (see also [20])

$$
\begin{aligned}
\vec{k}\cdot\vec{\varepsilon}^{\,*}_{\lambda'} &= \vec{k}\cdot\vec{\varepsilon}_{\lambda'}=0\ ,\ \vec{\varepsilon}^{\,*}_{\lambda'}\cdot\vec{\varepsilon}_{\lambda''}=\delta_{\lambda'\lambda''},\\
\sum_{\lambda'=1,2}\vec{\varepsilon}^{\,i*}_{\lambda'}\vec{\varepsilon}^{\,j}_{\lambda'} &= \delta^{ij}-\frac{\vec{k}^{\,i}\vec{k}^{\,j}}{\omega^2}=\delta^{ij}-\vec{n}^{\,i}\vec{n}^{\,j}, \qquad \text{(A-29)}
\end{aligned}
$$

where $\vec{n}=\vec{k}/\omega$. For the summation over only physical degrees of freedom of a real photon it is convenient to remove from Eq.(A-21) all terms proportional to the time–component of the photon polarization vector. This gives

$$
\begin{aligned}
&\sum_{\mathrm{pol}}\frac{|\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}|^2}{4m_n^2}=\frac{1}{(k_e\cdot k)^2}\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{(k_e\cdot k)^2}\Big[\lambda^2\Big(1-\frac{E_e+E_\nu+\omega}{M}\Big)\,\delta^{ij}-\frac{\lambda\,(\kappa+1)}{M}\,i\,\varepsilon^{ij\ell}(\vec{k}_e+\vec{k}_\nu+\vec{k}\,)^\ell\Big]\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{(k_e\cdot k)^2}\,\frac{\lambda^2+1}{2M}\,(\vec{k}_e+\vec{k}_\nu+\vec{k}\,)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\vec{\gamma}\,\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{(k_e\cdot k)^2}\,\frac{\lambda^2+1}{2M}\,(\vec{k}_e+\vec{k}_\nu+\vec{k}\,)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}\,\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{k_e\cdot k}\Big(-\frac{\lambda}{M}\,\vec{\varepsilon}^{\,*}_{\lambda'}+\frac{\lambda}{M\omega}\,i\,(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)\Big)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}\,\hat{k}_\nu\gamma^0\,\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{k_e\cdot k}\Big(-\frac{\lambda}{M}\,\vec{\varepsilon}_{\lambda'}-\frac{\lambda}{M\omega}\,i\,(\vec{\varepsilon}_{\lambda'}\times\vec{k}\,)\Big)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\vec{\gamma}\,(1-\gamma^5)\}
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{1}{k_e\cdot k}\Big(-\frac{\lambda^2}{M}\,\vec{\varepsilon}^{\,*}_{\lambda'}+i\,\frac{\lambda}{M\omega}\,(\vec{\varepsilon}^{\,*}_{\lambda'}\times\vec{k}\,)\Big)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\,\hat{k}_\nu\vec{\gamma}\,\bar{Q}_e(1-\gamma^5)\}\\
&+\frac{1}{k_e\cdot k}\Big(-\frac{\lambda^2}{M}\,\vec{\varepsilon}_{\lambda'}-\frac{\lambda}{M\omega}\,i\,(\vec{\varepsilon}_{\lambda'}\times\vec{k}\,)\Big)\cdot\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}\,\hat{k}_\nu\gamma^0(1-\gamma^5)\}\\
&+\frac{1}{k_e\cdot k}\Big[\frac{\lambda^2}{M\omega}\Big(\vec{\varepsilon}^{\,i*}_{\lambda'}\vec{k}^{\,j}-\vec{\varepsilon}^{\,j*}_{\lambda'}\vec{k}^{\,i}\Big)+\frac{\lambda^2}{M}\,i\,\varepsilon^{ij\ell}\vec{\varepsilon}^{\,\ell*}_{\lambda'}\Big]\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}\\
&-\frac{1}{k_e\cdot k}\Big[\frac{\lambda^2}{M\omega}\Big(\vec{\varepsilon}^{\,i}_{\lambda'}\vec{k}^{\,j}-\vec{\varepsilon}^{\,j}_{\lambda'}\vec{k}^{\,i}\Big)-\frac{\lambda^2}{M}\,i\,\varepsilon^{ij\ell}\,\vec{\varepsilon}^{\,\ell}_{\lambda'}\Big]\,\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,(1-\gamma^5)\}. \qquad \text{(A-30)}
\end{aligned}
$$

Summing over the photon polarizations we obtain

$$
\begin{aligned}
&\frac{1}{16}\sum_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\hat{k}_\nu\gamma^0\bar{Q}_e(1-\gamma^5)\}=E_eE_\nu\Big\{\Big[\Big(1+\frac{\omega}{E_e}\Big)\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\\
&+\frac{\vec{k}_e\cdot\vec{k}_\nu}{E_eE_\nu}\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\omega\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu\cdot\vec{n}}{E_\nu}\Big[\omega\Big(1+\frac{\omega}{E_e}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)-\omega\frac{m_e^2}{E_e}\Big]\Big\},\\
&\frac{1}{16}\sum_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}=E_\nu E_e\Big\{\delta^{ij}\Big[\Big(1+\frac{\omega}{E_e}\Big)\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\\
&-\delta^{ij}\Big\{\frac{\vec{k}_e\cdot\vec{k}_\nu}{E_eE_\nu}\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\omega\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu\cdot\vec{n}}{E_\nu}\Big[\omega\Big(1+\frac{\omega}{E_e}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)-\omega\frac{m_e^2}{E_e}\Big\}\\
&-i\,\varepsilon^{ij\ell}\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{(\vec{k}_e+\omega\vec{n}\,)^\ell}{E_e}+\frac{\omega}{E_e}\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\frac{\omega^2}{E_e}\,\vec{n}^{\,\ell}\,\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\\
&-i\,\varepsilon^{ij\ell}\frac{\vec{k}_\nu^{\,\ell}}{E_\nu}\Big[\Big(1+\frac{\omega}{E_e}\Big)\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu^{\,i}}{E_\nu}\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{(\vec{k}_e+\omega\vec{n}\,)^j}{E_e}\\
&+\frac{\omega}{E_e}\Big(\vec{k}_e^{\,j}-\vec{n}^{\,j}(\vec{k}_e\cdot\vec{n}\,)\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\frac{\omega^2}{E_e}\,\vec{n}^{\,j}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu^{\,j}}{E_\nu}\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{(\vec{k}_e+\omega\vec{n}\,)^i}{E_e}\\
&+\frac{\omega}{E_e}\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\frac{\omega^2}{E_e}\,\vec{n}^{\,i}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\Big\}\\
&\frac{1}{16}\sum_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\vec{\gamma}\,\hat{k}_\nu\,\gamma^0\,\bar{Q}_e(1-\gamma^5)\}=E_\nu E_e\Big\{\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{(\vec{k}_e+\omega\vec{n}\,)}{E_e}+\frac{\omega}{E_e}\Big(\vec{k}_e-\vec{n}\,(\vec{k}_e\cdot\vec{n}\,)\Big)\\
&\times\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\vec{n}\,\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu}{E_\nu}\Big[\Big(1+\frac{\omega}{E_e}\Big)\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\\
&+i\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{\vec{k}_\nu\times(\vec{k}_e+\omega\vec{n}\,)}{E_\nu E_e}+\frac{\omega^2}{E_e}\frac{\vec{k}_\nu\times\vec{n}}{E_\nu}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\omega\Big(\frac{\vec{k}_\nu\times\vec{k}_e}{E_\nu E_e}-\frac{(\vec{k}_\nu\times\vec{n}\,)(\vec{k}_e\cdot\vec{n}\,)}{E_\nu E_e}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big\},\\
&\frac{1}{16}\sum_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\,\hat{k}_\nu\,\vec{\gamma}\,\bar{Q}_e(1-\gamma^5)\}=E_\nu E_e\Big\{\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{(\vec{k}_e+\omega\vec{n}\,)}{E_e}+\frac{\omega}{E_e}\Big(\vec{k}_e-\vec{n}\,(\vec{k}_e\cdot\vec{n}\,)\Big)\\
&\times\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\vec{n}\,\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]+\frac{\vec{k}_\nu}{E_\nu}\Big[\Big(1+\frac{\omega}{E_e}\Big)\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)+\frac{\omega^2}{E_e}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big]\\
&-i\Big[\Big(k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2\Big)\frac{\vec{k}_\nu\times(\vec{k}_e+\omega\vec{n}\,)}{E_\nu E_e}+\frac{\omega^2}{E_e}\frac{\vec{k}_\nu\times\vec{n}}{E_\nu}\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)+\omega\Big(\frac{\vec{k}_\nu\times\vec{k}_e}{E_\nu E_e}-\frac{(\vec{k}_\nu\times\vec{n}\,)(\vec{k}_e\cdot\vec{n}\,)}{E_\nu E_e}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)\Big\},\\
&\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,i*}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,j}\,\hat{k}_\nu\gamma^0\,\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^j-\Big(\delta^{ij}-\vec{n}^{\,i}\vec{n}^{\,j}\Big)\\
&\times\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)-\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{j\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}-\omega\,i\,\varepsilon^{ij\ell}\,\vec{k}_e^{\,\ell}+\omega\,E_e\,i\,\varepsilon^{ij\ell}\,\vec{n}^{\,\ell}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)\\
&\times(2E_e+\omega)-\omega\,i\,\varepsilon^{i\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}\Big]-i\,\varepsilon^{ja\ell}\,\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^\ell-\Big(\delta^{i\ell}-\vec{n}^{\,i}\,\vec{n}^{\,\ell}\Big)\\
&\times\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\,\Big)-\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{\ell bc}\,\vec{n}^{\,b}\,\vec{k}_e^{\,c}+\omega\,i\,\varepsilon^{\ell ib}\vec{k}_e^{\,b}-\omega\,E_e\,i\,\varepsilon^{\ell ia}\,\vec{n}^{\,a}\Big],\\
&\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,i}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\gamma^0\,\hat{k}_\nu\vec{\gamma}^{\,j}\,(1-\gamma^5)\}=E_\nu\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^j-\Big(\delta^{ij}-\vec{n}^{\,i}\vec{n}^{\,j}\Big)
\end{aligned}
$$

$$\times\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{j\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}+\omega\,i\,\varepsilon^{ij\ell}\,\vec{k}_e^{\,\ell}-\omega\,E_e\,i\,\varepsilon^{ij\ell}\,\vec{n}^{\,\ell}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)$$
$$\times\,(2E_e+\omega)+\omega\,i\,\varepsilon^{i\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}\Big]+i\,\varepsilon^{ja\ell}\,\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^\ell-\Big(\delta^{i\ell}-\vec{n}^{\,i}\,\vec{n}^{\,\ell}\Big)$$
$$\times\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{\ell bc}\,\vec{n}^{\,b}\,\vec{k}_e^{\,c}-\omega\,i\,\varepsilon^{\ell ib}\,\vec{k}_e^{\,b}+\omega\,E_e\,i\,\varepsilon^{\ell ia}\,\vec{n}^{\,a}\Big],$$

$$\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,i*}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,\gamma^0\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}=E_\nu\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^j-\Big(\delta^{ij}-\vec{n}^{\,i}\vec{n}^{\,j}\Big)$$
$$\times\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)-\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{j\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}-\omega\,i\,\varepsilon^{ij\ell}\,\vec{k}_e^{\,\ell}+\omega\,E_e\,i\,\varepsilon^{ij\ell}\,\vec{n}^{\,\ell}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)$$
$$-\omega\,i\,\varepsilon^{j\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}\Big]-i\,\varepsilon^{ja\ell}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^\ell-\Big(\delta^{i\ell}-\vec{n}^{\,i}\vec{n}^{\,\ell}\Big)\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)-\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{\ell ab}\,\vec{n}^{\,a}\,\vec{k}_e^{\,b}$$
$$+\omega\,i\,\varepsilon^{\ell ib}\,\vec{k}_e^{\,b}-\omega\,E_e\,i\,\varepsilon^{\ell ib}\,\vec{n}^{\,b}\Big],$$

$$\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,i}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\,\vec{\gamma}^{\,j}\,\hat{k}_\nu\gamma^0\,(1-\gamma^5)\}=E_\nu\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^j-\Big(\delta^{ij}-\vec{n}^{\,i}\vec{n}^{\,j}\Big)$$
$$\times\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{j\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}+\omega\,i\,\varepsilon^{ij\ell}\,\vec{k}_e^{\,\ell}-\omega\,E_e\,i\,\varepsilon^{ij\ell}\,\vec{n}^{\,\ell}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)$$
$$+\omega\,i\,\varepsilon^{i\ell a}\,\vec{n}^{\,\ell}\,\vec{k}_e^{\,a}\Big]-i\,\varepsilon^{ja\ell}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,i}-\vec{n}^{\,i}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^\ell-\Big(\delta^{i\ell}-\vec{n}^{\,i}\vec{n}^{\,\ell}\Big)\,\omega\,\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,i}\,i\,\varepsilon^{\ell ab}\,\vec{n}^{\,a}\,\vec{k}_e^{\,b}$$
$$-\omega\,i\,\varepsilon^{\ell ib}\,\vec{k}_e^{\,b}+\omega\,E_e\,i\,\varepsilon^{\ell ib}\,\vec{n}^{\,b}\Big],$$

$$\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,\ell*}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,\bar{Q}_e(1-\gamma^5)\}=\delta^{ij}E_\nu\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)-\omega\,i\,\varepsilon^{\ell bc}\vec{n}^{\,b}\vec{k}_e^{\,c}\Big]$$
$$-i\,\varepsilon^{ija}E_\nu\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,a}-\omega\Big(\delta^{\ell a}-\vec{n}^{\,\ell}\vec{n}^{\,a}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)-\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{abc}\vec{n}^{\,b}\vec{k}_e^{\,c}+\omega\,i\,\varepsilon^{a\ell b}\vec{k}_e^{\,b}$$
$$-\omega E_e\,i\,\varepsilon^{a\ell b}\vec{n}^{\,b}\Big]-\delta^{ij}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,a}-\omega\Big(\delta^{\ell a}-\vec{n}^{\,\ell}\vec{n}^{\,a}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)-\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{abc}\vec{n}^{\,b}\vec{k}_e^{\,c}$$
$$+\omega\,i\,\varepsilon^{a\ell b}\vec{k}_e^{\,b}-\omega E_e\,i\,\varepsilon^{a\ell b}\vec{n}^{\,b}\Big]+\vec{k}_\nu^{\,i}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,j}-\omega\Big(\delta^{\ell j}-\vec{n}^{\,\ell}\vec{n}^{\,j}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)$$
$$-\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{jbc}\vec{n}^{\,b}\vec{k}_e^{\,c}+\omega\,i\,\varepsilon^{j\ell b}\vec{k}_e^{\,b}-\omega E_e\,i\,\varepsilon^{j\ell b}\vec{n}^{\,b}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,i}-\omega\Big(\delta^{\ell i}-\vec{n}^{\,\ell}\vec{n}^{\,i}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)$$
$$-\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{ibc}\vec{n}^{\,b}\vec{k}_e^{\,c}+\omega\,i\,\varepsilon^{i\ell b}\vec{k}_e^{\,b}-\omega E_e\,i\,\varepsilon^{i\ell b}\vec{n}^{\,b}\Big]-i\,\varepsilon^{ija}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)-\omega\,i\,\varepsilon^{\ell bc}\vec{n}^{\,b}\vec{k}_e^{\,c}\Big],$$

$$\frac{1}{4}\sum_{\lambda'}\vec{\varepsilon}^{\,\ell}_{\lambda'}\mathrm{tr}\{(\hat{k}_e+m_e)\,Q_e\,\vec{\gamma}^{\,i}\,\hat{k}_\nu\vec{\gamma}^{\,j}\,(1-\gamma^5)\}=\delta^{ij}E_\nu\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)+\omega\,i\,\varepsilon^{\ell bc}\vec{n}^{\,b}\vec{k}_e^{\,c}\Big]$$
$$-i\,\varepsilon^{ija}E_\nu\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,a}-\omega\Big(\delta^{\ell a}-\vec{n}^{\,\ell}\vec{n}^{\,a}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{abc}\vec{n}^{\,b}\vec{k}_e^{\,c}-\omega\,i\,\varepsilon^{a\ell b}\vec{k}_e^{\,b}$$
$$+\omega E_e\,i\,\varepsilon^{a\ell b}\vec{n}^{\,b}\Big]-\delta^{ij}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,a}-\omega\Big(\delta^{\ell a}-\vec{n}^{\,\ell}\vec{n}^{\,a}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)+\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{abc}\vec{n}^{\,b}\vec{k}_e^{\,c}$$
$$-\omega\,i\,\varepsilon^{a\ell b}\vec{k}_e^{\,b}+\omega E_e\,i\,\varepsilon^{a\ell b}\vec{n}^{\,b}\Big]+\vec{k}_\nu^{\,i}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,j}-\omega\Big(\delta^{\ell j}-\vec{n}^{\,\ell}\vec{n}^{\,j}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)$$
$$+\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{jbc}\vec{n}^{\,b}\vec{k}_e^{\,c}-\omega\,i\,\varepsilon^{j\ell b}\vec{k}_e^{\,b}+\omega E_e\,i\,\varepsilon^{j\ell b}\vec{n}^{\,b}\Big]+\vec{k}_\nu^{\,j}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2\vec{k}_e+\omega\vec{n}\,)^{\,i}-\omega\Big(\delta^{\ell i}-\vec{n}^{\,\ell}\vec{n}^{\,i}\Big)\Big(E_e-\vec{k}_e\cdot\vec{n}\Big)$$
$$+\omega\,\vec{n}^{\,\ell}\,i\,\varepsilon^{ibc}\vec{n}^{\,b}\vec{k}_e^{\,c}-\omega\,i\,\varepsilon^{i\ell b}\vec{k}_e^{\,b}+\omega E_e\,i\,\varepsilon^{i\ell b}\vec{n}^{\,b}\Big]-i\,\varepsilon^{ija}\vec{k}_\nu^{\,a}\Big[-\Big(\vec{k}_e^{\,\ell}-\vec{n}^{\,\ell}(\vec{k}_e\cdot\vec{n}\,)\Big)(2E_e+\omega)+\omega\,i\,\varepsilon^{\ell bc}\vec{n}^{\,b}\vec{k}_e^{\,c}\Big]. \qquad \text{(A-31)}$$

The rate of the neutron radiative β^-–decay is given by

$$\lambda_{\beta\gamma} \;=\; \pi\alpha G_F^2|V_{ud}|^2\frac{1}{m_n}\int\frac{1}{2}\sum_{\lambda',\mathrm{pol}}|\mathcal{M}(n\to pe^-\bar{\nu}_e\gamma)_{\lambda'}|^2F(E_e,Z=1)$$
$$\times\,(2\pi)^4\delta^{(4)}(k_n-k_p-k_e-k_\nu-k)\,\frac{d^3k_p}{(2\pi)^3 2E_p}\frac{d^3k_e}{(2\pi)^3 2E_e}\frac{d^3k_\nu}{(2\pi)^3 2E_\nu}\frac{d^3k}{(2\pi)^3 2\omega}, \qquad \text{(A-32)}$$

where $F(E_e,Z=1)$ is the relativistic Fermi function, taking into account the final–state proton–electron Coulomb interaction [10, 11] (see also Eq.(8)). After the integration over the proton momentum $\vec{k}_p$, the directions of the

momenta of the particles in the final state we are left with the result [10]

$$\lambda_{\beta\gamma}(\omega_{\rm max},\omega_{\rm min}) = (1+3\lambda^2)\frac{\alpha}{\pi}\frac{G_F^2|V_{ud}|^2}{2\pi^3}\int_{\omega_{\rm min}}^{\omega_{\rm max}}\frac{d\omega}{\omega}\int_{m_e}^{E_0-\omega} dE_e\, E_e\sqrt{E_e^2-m_e^2}(E_0-E_e-\omega)^2\, F(E_e,Z=1)\,\rho_{\beta\gamma}(E_e,\omega), \tag{A-33}$$

where the rate of the neutron radiative β^-–decay is defined for the photon energy region $\omega_{\rm min}\leq\omega\leq\omega_{\rm max}$ and $E_0=(m_n^2-m_p^2+m_e^2)/2m_n$ is the end–point energy of the electron energy–spectrum of the neutron β^-–decay [10]. The function $\rho_{\beta\gamma}(E_e,\omega)$ is given by

$$\rho_{\beta\gamma}(E_e,\omega) = \int\frac{d\Omega_{e\gamma}}{4\pi}\frac{d\Omega_\nu}{4\pi}\,\frac{1}{2}\sum_{\lambda',{\rm pol}}\frac{\omega^2|\mathcal{M}(n\to pe^-\bar\nu_e\gamma)_{\lambda'}|^2}{(1+3\lambda^2)32m_n^2E_eE_\nu}\,\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu), \tag{A-34}$$

where $d\Omega_{e\gamma}=\sin\theta_{e\gamma}d\theta_{e\gamma}d\varphi_{e\gamma}$ is an infinitesimal solid angle of the electron–photon momentum correlations $\vec k_e\cdot\vec n = k_e\cos\theta_{e\gamma}$, defined by the polar angle $0\leq\theta_{e\gamma}\leq\pi$ and the azimuthal angle $0\leq\varphi_{e\gamma}\leq 2\pi$, and $d\Omega_\nu$ is an infinitesimal solid angle of the antineutrino momentum $\vec k_\nu$. The function $\Phi_{\beta\gamma}(\vec k_e,\vec k\,\vec k_\nu)$ is defined by the integral [10, 11].

$$\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu) = \int_0^\infty\delta(m_n-\sqrt{m_p^2+(\vec k_e+\vec k+\vec k_\nu)^2}-E_e-E_\nu-\omega)\,\frac{m_n}{E_p}\,\frac{E_\nu^2dE_\nu}{(E_0-E_e-\omega)^2}. \tag{A-35}$$

The result of the integration over E_ν is equal to

$$\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu) = \frac{\Big(1+\dfrac{1}{E_0-E_e-\omega}\dfrac{k_e\cdot k}{m_n}\Big)^2}{\Big(1-\dfrac{1}{m_n}\dfrac{(k_e+k)\cdot k_\nu}{E_\nu}\Big)^3}, \tag{A-36}$$

where the antineutrino energy is defined by

$$E_\nu = \frac{E_0-E_e-\omega+\dfrac{k_e\cdot k}{m_n}}{1-\dfrac{1}{m_n}\,(E_e+\omega-|\vec k_e+\vec k\,|\cos\theta_\nu)}. \tag{A-37}$$

Here θ_ν is an angle between the momenta $\vec k_e+\vec k$ and $\vec k_\nu$. To order $1/M$ the function $\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu)$ is given by

$$\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu) = 1+\frac{2}{M}\,\frac{\omega}{E_\nu}\Big(E_e-\vec k_e\cdot\vec n\,\Big)+\frac{3}{M}\Big(E_e+\omega-\frac{(\vec k_e+\omega\vec n\,)\cdot\vec k_\nu}{E_\nu}\Big), \tag{A-38}$$

where $E_\nu=E_0-E_e-\omega$ [10]. Setting $\omega=0$ we arrive at the corresponding function of the neutron β^-–decay [10]. For the calculation of the function $\rho_{\beta\gamma}(E_e,\omega)$ we propose, first, to integrate over the directions of the antineutrino 3–momentum. Now making the integration in Eq.(A-34) over the antineutrino momentum solid angle we arrive at the expression

$$\begin{aligned}
&\int\frac{d\Omega_\nu}{4\pi}\,\frac{1}{2}\sum_{\lambda',{\rm pol}}\frac{\omega^2|\mathcal{M}(n\to pe^-\bar\nu_e\gamma)_{\lambda'}|^2}{(1+3\lambda^2)32m_n^2E_eE_\nu}\,\Phi_{\beta\gamma}(\vec k_e,\vec k,\vec k_\nu) = \Big[1+2\,\frac{\omega}{M}\,\frac{E_e-\vec k_e\cdot\vec n}{E_0-E_e-\omega}+\frac{3}{M}\,\Big(E_e+\omega-\frac{1}{3}\,E_0\Big)\\
&+\frac{\lambda^2-2(\kappa+1)\lambda+1}{1+3\lambda^2}\,\frac{E_0-E_e-\omega}{M}\Big]\Big[\Big(1+\frac{\omega}{E_e}\Big)\,\frac{k_e^2-(\vec k_e\cdot\vec n\,)^2}{(E_e-\vec k_e\cdot\vec n\,)^2}+\frac{\omega^2}{E_e}\,\frac{1}{E_e-\vec k_e\cdot\vec n}\Big]+\frac{3\lambda^2-1}{1+3\lambda^2}\,\frac{1}{M}\\
&\times\Big\{\frac{k_e^2+\omega\vec k_e\cdot\vec n}{E_e}\,\Big(\frac{k_e^2-(\vec k_e\cdot\vec n\,)^2}{(E_e-\vec k_e\cdot\vec n\,)^2}+\frac{\omega}{E_e-\vec k_e\cdot\vec n}\Big)+(\omega+\vec k_e\cdot\vec n\,)\Big[\Big(1+\frac{\omega}{E_e}\Big)\frac{\omega}{E_e-\vec k_e\cdot\vec n}-\frac{m_e^2}{E_e}\,\frac{\omega}{(E_e-\vec k_e\cdot\vec n\,)^2}\Big]\Big\}\\
&-\frac{\lambda^2+2(\kappa+1)\lambda-1}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{k_e^2+\omega^2+2\omega\vec k_e\cdot\vec n}{E_e}\,\frac{k_e^2-(\vec k_e\cdot\vec n\,)^2}{(E_e-\vec k_e\cdot\vec n\,)^2}+\frac{\omega}{E_e}\,\frac{k_e^2-(\vec k_e\cdot\vec n\,)^2}{E_e-\vec k_e\cdot\vec n}+\frac{\omega^2}{E_e}\,\frac{\omega+\vec k_e\cdot\vec n}{E_e-\vec k_e\cdot\vec n}\Big]\\
&-\frac{\lambda(\lambda-1)}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{\omega}{E_e}\,\frac{k_e^2-(\vec k_e\cdot\vec n\,)^2}{E_e-\vec k_e\cdot\vec n}+3\,\frac{\omega^2}{E_e}\Big].
\end{aligned} \tag{A-39}$$

Plugging Eq.(A-39) into Eq.(A-34) we define the function $\rho_{\beta\gamma}(E_e,\omega)$ by the integral over electron–photon correlation angles only. It is given by

$$\rho_{\beta\gamma}(E_e,\omega) = \int \frac{d\Omega_{e\gamma}}{4\pi}\Big[1+2\,\frac{\omega}{M}\,\frac{E_e-\vec{k}_e\cdot\vec{n}}{E_0-E_e-\omega}+\frac{3}{M}\Big(E_e+\omega-\frac{1}{3}\,E_0\Big)+\frac{\lambda^2-2(\kappa+1)\lambda+1}{1+3\lambda^2}\,\frac{E_0-E_e-\omega}{M}\Big]$$
$$\times\Big[\Big(1+\frac{\omega}{E_e}\Big)\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega^2}{E_e}\,\frac{1}{E_e-\vec{k}_e\cdot\vec{n}}\Big]+\frac{3\lambda^2-1}{1+3\lambda^2}\,\frac{1}{M}\,\Big\{\frac{k_e^2+\omega\vec{k}_e\cdot\vec{n}}{E_e}\,\Big(\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega}{E_e-\vec{k}_e\cdot\vec{n}}\Big)$$
$$+(\omega+\vec{k}_e\cdot\vec{n}\,)\Big[\Big(1+\frac{\omega}{E_e}\Big)\frac{\omega}{E_e-\vec{k}_e\cdot\vec{n}}-\frac{m_e^2}{E_e}\,\frac{\omega}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}\Big]\Big\}-\frac{\lambda^2+2(\kappa+1)\lambda-1}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{k_e^2+\omega^2+2\omega\vec{k}_e\cdot\vec{n}}{E_e}$$
$$\times\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{(E_e-\vec{k}_e\cdot\vec{n}\,)^2}+\frac{\omega}{E_e}\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{E_e-\vec{k}_e\cdot\vec{n}}+\frac{\omega^2}{E_e}\,\frac{\omega+\vec{k}_e\cdot\vec{n}}{E_e-\vec{k}_e\cdot\vec{n}}\Big]-\frac{\lambda(\lambda-1)}{1+3\lambda^2}\,\frac{1}{M}\,\Big[\frac{\omega}{E_e}\,\frac{k_e^2-(\vec{k}_e\cdot\vec{n}\,)^2}{E_e-\vec{k}_e\cdot\vec{n}}+3\,\frac{\omega^2}{E_e}\Big]. \qquad \text{(A-40)}$$

To leading order in large proton mass expansion, i.e. at $M\to\infty$, Eq.(A-40) reduces to the expression, calculated in [10].

[1] S. M. Berman, Phys. Rev. **112**, 267 (1958).
[2] T. Kinoshita and A. Sirlin, Phys. Rev. **113**, 1652 (1959).
[3] S. M. Berman and A. Sirlin, Ann. Phys. (N.Y.) **20**, 20 (1962).
[4] G. Källén, Nucl. Phys. B **1**, 225 (1967).
[5] A. Sirlin, Phys. Rev. **164**, 1767 (1967).
[6] E. S. Abers, D. A. Dicus, R. E. Norton, and H. R. Queen, Phys. Rev. **167**, 1461 (1968).
[7] R. T. Shann, Nuovo Cimento A **5**, 591 (1971).
[8] Yu. V. Gaponov and R. U. Khafizov, Phys. At. Nucl. **59**, 1213 (1996) (Yad. Fiz. **59**, 1270 (1996)); Phys. Lett. B **379**,7 (1996); Nucl. Instrum. Methods Phys. Res., Sect. A **440**, 557 (2000).
[9] V. Bernard, S. Gardner, U.-G. Meißner, and C. Zang, Phys. Lett. B **593**, 105 (2004); Erratum Phys. Lett. B **599**, 348 (2004).
[10] A. N. Ivanov, M. Pitschmann, and N. I. Troitskaya, Phys. Rev. D **88**, 073002 (2013); arXiv:1212.0332v4 [hep–ph].
[11] A.N. Ivanov, R. Höllwieser, N. I. Troitskaya, and M. Wellenzohn, Phys. Rev. D **88**, 065026 (2013).
[12] J. S. Nico *et al.*, Nature, **444**, 1059 (2006).
[13] R. L. Cooper *et al.*, Phys. Rev. C **81**, 035503 (2010).
[14] S. Gardner and D. He, Phys. Rev. D **86**, 016003 (2012); Hyperfine Interact. **214**, 71 (2013).
[15] S. Gardner and D. He, Phys. Rev. D **87**, 116012 (2013).
[16] M. J. Bales *et al.* (RDK II Collaboration), Phys. Rev. Lett. **116**, 242501 (2016).
[17] H. Abele, Progr. Part. Nucl. Phys. **60**, 1 (2008).
[18] C. Partignani *et al.* (Particle Data Group), Chinese Physics C **40**, 100001 (2016).
[19] C. Itzykson and J.–B. Zuber, in *Quantum Field Theory*, McGraw–Hill Inc., New York, 1980.
[20] A. N. Ivanov, R. Höllwieser, N. I. Troitskaya, M. Wellenzohn, O. M. Zherebtsov, and A. P. Serebrov, Phys. Rev. C **88**, 055501 (2013).
[21] J. D. Bjorken and S. D. Drell, in *Relativistische Quantenfeldtheorie*, Bibliographisches Institut Mannheim, B•I•, Hochschultaschenbücher - Verlag, 1967.
[22] R. Pohl *et al.*, Nature **466**, 213 (2010).
[23] A. N. Ivanov, M. Pitschmann, N. I. Troitskaya, and Ya. A. Berdnikov, Phys. Rev. C **89**, 055502 (2014).